

Relic density in nonuniversal gaugino mass models with SO(10) GUT symmetry

Katri Huitu*

*Department of Physics, and Helsinki Institute of Physics,
P.O. Box 64, FIN-00014 University of Helsinki, Finland*

Jari Laamanen†

*Institut für Physik, Technische Universität Dortmund, D-44221 Dortmund, Germany
(Dated: April 11, 2009)*

Nonuniversal boundary conditions in grand unified theories can lead to nonuniversal gaugino masses at the unification scale. In R -parity preserving theories the lightest supersymmetric particle is a natural candidate for the dark matter. The composition of the lightest neutralino and the identity of the next-to-lightest supersymmetric particle are studied, when nonuniversal gaugino masses come from representations of SO(10). In these cases, the thermal relic density compatible with the Wilkinson Microwave Anisotropy Probe observations is found. Relic densities are compared with the universal case. Mass spectra in the studied cases are discussed.

PACS numbers: 12.60.Jv, 95.35.+d

Keywords: Gaugino masses, relic density, dark matter

I. INTRODUCTION

The phenomenology of supersymmetric models depends crucially on the compositions of neutralinos and charginos, if the lightest neutralino is the lightest supersymmetric particle (LSP). In addition to the laboratory studies, relevant input is obtained from the dark matter searches, where the Wilkinson Microwave Anisotropy Probe (WMAP) satellite has put precise limits on the relic density. Supersymmetric theories which preserve R -parity contain a natural candidate for the cold dark matter particle. Neutralino LSP can provide the appropriate relic density.

In many supergravity type models the lightest neutralino is binolike, which often leads to too large thermal relic density, as compared to the limits provided by the WMAP experiment [1]. When the gaugino masses are not universal at the grand unification scale, the resulting neutralino composition changes from the case of universal gaugino masses [2]. In this paper, the thermal relic density of the neutralino LSP is studied, when gaugino masses are due to nonuniversal representations of SO(10) grand unified theory (GUT) [3, 4]. Dark matter in a particular gauge symmetry breaking chain of the SO(10) GUT in the case of universal gaugino masses has been recently studied in [5]. Some phenomenological aspects of SO(10) GUTs with nonuniversal gaugino masses have been considered in [6, 7].

SO(10) has many attractive features among the GUT models. One of the most appealing properties is that one family of matter fermions can be put into a single 16-dimensional irreducible spinor representation of SO(10), including the right-handed neutrino [8, 9]. In

addition, SO(10) allows possibility for the Yukawa coupling unification and representations are anomaly free. Conservation of R -parity, which forbids the unwanted dimension-five operators leading to rapid proton decay, may result from the SO(10) symmetry breaking. The doublet-triplet splitting could be achieved using, *e.g.*, the so-called Dimopoulos-Wilczek mechanism [10]. Because the SO(10) gauge symmetry breaks down to the standard model (SM) gauge symmetry through some intermediate group, the SO(10) GUT offers several possibilities for the model building. For example, it can contain as a subgroup the Pati-Salam $SU(4) \times SU(2) \times SU(2)$ model.

Gaugino masses originate from the non-renormalizable terms in the $N = 1$ supergravity Lagrangian involving the gauge kinetic function $f_{ab}(\Phi)$ [11]. The gauge part of the Lagrangian contains the gauge kinetic function coupling with two field strength superfields W^a . The Lagrangian for the coupling can be written as

$$\mathcal{L}_{gk} = \int d^2\theta f_{ab}(\Phi) W^a W^b + \text{H.C.}, \quad (1)$$

where a and b are gauge group indices (for example, $a, b = 1, 2, \dots, 45$ for SO(10)), and repeated indices are summed over. The function $f_{ab}(\Phi)$ is an analytic function of the chiral superfields Φ in the theory. The chiral superfields Φ consist of a set of gauge singlet superfields Φ^s and gauge nonsinglet superfields Φ^n under the grand unified group. The gauge kinetic function $f_{ab}(\Phi)$ can be expanded,

$$f_{ab}(\Phi) = f_0(\Phi^s) \delta_{ab} + \sum_n f_n(\Phi^s) \frac{\Phi_{ab}^n}{M_P} + \dots, \quad (2)$$

where Φ^s and Φ^n are the singlet and nonsinglet chiral superfields, respectively. Here $f_0(\Phi^s)$ and $f_n(\Phi^s)$ are functions of gauge singlet superfields Φ^s , and M_P is some large scale. In order to generate a mass term for the gauginos, the gauge kinetic function must be non-minimal,

*Electronic address: katri.huitu@helsinki.fi

†Electronic address: jari.laamanen@uni-dortmund.de

i.e., it must not be a constant [12]. When F_Φ gets a vacuum expectation value (vev) $\langle F_\Phi \rangle$, the interaction (1) gives rise to gaugino masses:

$$\mathcal{L}_{gk} \supset \frac{\langle F_\Phi \rangle_{ab}}{M_P} \lambda^a \lambda^b + \text{H.C.}, \quad (3)$$

where $\lambda^{a,b}$ are gaugino fields. The nonuniversal gaugino masses are generated by the nonsinglet chiral superfield Φ^n that appears linearly in the gauge kinetic function $f_{ab}(\Phi)$ in Eq. (2).

Gauginos belong to the adjoint representation of the gauge group, which in the case of $\text{SO}(10)$ is the **45** dimensional representation. Because Eq. (3) must be gauge invariant, Φ and F_Φ must belong to some of the following representations appearing in the symmetric product of the two **45** dimensional representations of $\text{SO}(10)$ [13, 14]:

$$(\mathbf{45} \otimes \mathbf{45})_{\text{Symm}} = \mathbf{1} \oplus \mathbf{54} \oplus \mathbf{210} \oplus \mathbf{770}. \quad (4)$$

The representations **54**, **210** and **770** may lead to nonuniversal gaugino masses, while the **1** dimensional representation gives manifestly the universal gaugino masses. The relations between the gaugino masses are determined by the representation invariants, and are specific for each of the representations. Because the gauge kinetic function in Eq. (2) can get contributions from several different Φ 's, a linear combination of any of the representations is also possible. In that case the gaugino mass terms are not uniquely determined anymore, in contrast to the contribution purely from one representation. Here we assume that the dominant component of the gaugino masses comes from only one representation. This gives us a clear understanding of the role of different representations.

II. DARK MATTER IN $\text{SO}(10)$ REPRESENTATIONS

A. Breaking Chains: $\text{SO}(10) \rightarrow H \rightarrow \text{SM}$

The GUT group $\text{SO}(10)$ breaks down to the standard model gauge group $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ via some intermediate gauge group H . Therefore the gaugino mass relations depend also on the gauge group breaking chain, in addition to the representation invariants coming from the gauge kinetic function. Moreover, the intermediate breaking scale affects also the generated gaugino masses via heavy gauge supermultiplets that correspond to the broken generators. However, if the gauge breaking from the GUT group to the SM group takes place at the GUT scale, these loop-induced messenger contributions [15] can be neglected in comparison to the tree-level contributions. Some fits to the experimental data in $\text{SO}(10)$ GUT indicate that the two breaking scales are very close to each other, see [16, 17], although realistic models exist also with large splitting of the scales [18]. In this

work we assume that the breaking from $\text{SO}(10)$ to the SM gauge group happens at the GUT scale, and that the GUT breaking does not affect the gauge coupling unification.

We will study the representations **54** and **210** in the right-hand side of Eq. (4). The interesting breaking chains of **54** and **210** are included also in the breaking chains of **770**. Table I shows possible $\text{SO}(10)$ breaking chains [14, 19, 20], which include the standard model gauge group, for the two chosen representations. Some of the subgroups lead to universal gaugino masses, or to massless gauginos [14], and we do not consider them. We will limit ourselves to the intermediate gauge groups $\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$, $\text{SU}(2) \times \text{SO}(7)$ and $\text{SU}(5) \times \text{U}(1)$.

TABLE I: Breaking chains of $\text{SO}(10)$ representations **54** and **210** which include the SM gauge group.

F_Φ	H	Subgroup description
54	$\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$	Pati-Salam
	$\text{SU}(2) \times \text{SO}(7)$	
	$\text{SO}(9)$	Universal gauginos
210	$\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$	Massless gluino
	$\text{SU}(3) \times \text{SU}(2) \times \text{SU}(2) \times \text{U}(1)$	Massless $\text{SU}(2)_L$ gauginos
	$\text{SU}(3) \times \text{SU}(2) \times \text{U}(1) \times \text{U}(1)$	
	$\text{SU}(5) \times \text{U}(1)$	“Flipped” $\text{SU}(5)$

Table II displays the ratios of resulting gaugino masses at the tree level as they arise when F_Φ belongs to the above-mentioned representations of $\text{SO}(10)$ or singlet [14]. The resulting 1-loop relations at the electroweak scale are also displayed. These values and the resulting

TABLE II: Ratios of the gaugino masses at the GUT scale in the normalization $M_3(\text{GUT}) = 1$, and at the electroweak scale in the normalization $M_3(\text{EW}) = 1$.

F_Φ	H	M_1^G	M_2^G	M_3^G	M_1^{EW}	M_2^{EW}	M_3^{EW}
1		1	1	1	0.14	0.29	1
54	$\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$	-1	-1.5	1	-0.15	-0.44	1
54	$\text{SU}(2) \times \text{SO}(7)$	1	-7/3	1	0.15	-0.68	1
210	$\text{SU}(5) \times \text{U}(1)$	-96/25	1	1	-0.56	0.29	1

relic densities can be compared with the universal and nonuniversal representations resulting in the $\text{SU}(5)$ GUT model [21, 22]. Since we assume breaking at one scale, the universal model with which we should compare in the $\text{SO}(10)$ GUT is similar to the universal model in the $\text{SU}(5)$ GUT. In the nonuniversal representations, the relations between gaugino masses change. Thus, *e.g.*, the **54**-dimensional Pati-Salam model of $\text{SO}(10)$ may seem at first glance rather similar to **24** of $\text{SU}(5)$, but we will see that the twice as large bino component has a large effect to the relic density. The bino and wino mass parameters affect directly the lightest neutralino mass and properties. They also affect the value of the μ -parameter through the renormalization group equations (RGE) and

the radiative electroweak symmetry breaking (rEWSB), therefore controlling also the Higgsino component in the lightest neutralino. Since the lightest neutralino mass limit can be deduced from the chargino mass limit, the nonuniversal gaugino masses change the lower limit for the neutralino mass: for **54** the neutralino mass limit is smaller than in the universal case, while for **210** the mass limit is close to the chargino mass limit.

B. Calculation of Dark Matter Relic Density

We calculate the SUSY spectrum for each model with the program SOFTSUSY (version 2.0.11) [23], and the resulting relic density with the program micrOMEGAs (version 2.0.7) [24, 25, 26]. For the relic density, we use here the WMAP combined 3 yr limits [1]

$$\Omega_{CDM}h^2 = 0.11054^{+0.00976}_{-0.00956} \quad (2\sigma). \quad (5)$$

In all the figures that we show below, the filling denoted by *wmap* is the WMAP-preferred region. For the $b \rightarrow s\gamma$ experimental branching fraction, we have used the two sigma world average [27],

$$BR(b \rightarrow s\gamma) = (355 \pm 24^{+9}_{-10} \pm 3) \times 10^{-6}. \quad (6)$$

The areas enclosed by the *bsg* contour are disallowed by the $b \rightarrow s\gamma$ constraint. For the particle masses, the following limits are applied [26]: $m_{\tilde{e}_R} > 99.4$ or 100.5 GeV depending on if the lightest neutralino mass is below or above 40 GeV, $m_{\tilde{\mu}_R} > 95$ GeV, $m_{\tilde{\tau}_1} > 80.5$ to 88 GeV depending on the lightest neutralino mass (from 10 to 75 GeV), $m_{\tilde{\nu}_i} > 43$ GeV, and $m_{\tilde{\chi}^\pm} > 73.1$ to 103 GeV depending on the sneutrino masses (from 45 to 425 GeV). In the figures, *lep* shows an area where the experimental mass limits are not met, *rge* shows an area where there is no radiative EWSB, and *lsp* the area where neutralino is not the LSP. The curve $m_h = 114$ GeV is depicted in the figures (dash-dotted line denoted by *h*). For the shown parameter regions, when otherwise experimentally allowed, Higgs is always heavier than 91 GeV, which is the Higgs mass limit in MSSM for $\tan\beta \geq 10$ assuming maximal top mixing [28].

1. Representation 54

The area of preferred thermal relic density for the two chains of the **54** dimensional representation are shown for sets of parameters in Figs. 1, 2, and 3. In each set of three figures, the first figure (a) represents the neutralino relic density for given parameters with collider constraints depicted in the plot, the second figure (b) shows, for the same parameters, the identity of the next-to-lightest supersymmetric particle, and the third figure (c) shows the lightest neutralino composition in RGB-color encoding, (*i.e.*, colors, or hues of black and white, indicate the particle as shown in the figure; therefore the mixture of the

colors, or hues of black and white, describes the nature of the $\tilde{\chi}_1^0$ -composition). In each of the figures, the WMAP-preferred relic density filling is also superimposed to the graph.

As can be seen from the Table II, the lightest neutralino is expected to be bino rather than wino. The large bino component tends to suppress the neutralino annihilation cross section, since bino lacks the s-channel Z-boson annihilation mode. A substantial Higgsino component is usually needed to help to increase the annihilation rate, unless there happens to be coannihilation or, *e.g.*, an open Higgs s-channel annihilation mode available.

In Fig. 1 relic density, the next-to-lightest supersymmetric particle (NLSP) and LSP composition in the breaking chain $SU(2) \times SO(7)$ are shown. Because $\tilde{\chi}_1^0$ is mostly bino, the spectrum with preferred relic density is quite light and conflicts with collider constraints in some parts of the parameter space. With increasing gaugino masses also the Higgsino component in the neutralino LSP increases, and at the point where the change to dominantly Higgsino LSP occurs, also the relic density drops. The overall relic density is not very high, thus allowing a wider WMAP-preferred region than, *e.g.*, in the singlet, *i.e.*, mSUGRA case [22]. For a given M_2 , the corresponding M_1 is smaller than in the singlet case, which results in a smaller μ value at the EW-scale. This has an effect of an increasing the Higgsino component in the lightest neutralino thus boosting the annihilation. The NLSP is chargino, and with increasing Higgsino component it eventually becomes the LSP.

In the allowed region, M_2 is less than 740 GeV, which restricts the lightest chargino mass to values less than ~ 150 GeV. The lower limit for the chargino mass is the LEP limit. The partners of the SM fermions are heavier than 300–500 GeV. Thus, assuming that the neutralino is responsible for the dark matter, this breaking chain of the gauge symmetry has as a robust prediction for the upper limit of the chargino mass, and furthermore it is lighter than the squarks and sleptons. For $M_2 \sim 350$ GeV the whole neutralino and chargino spectrum, and even the gluino, is lighter than the sfermions for the LEP allowed and WMAP-preferred region. The same is true also for $M_2 \sim 630$ GeV and $m_0 \gtrsim 940$ GeV for the WMAP-preferred region.

The $b \rightarrow s\gamma$ constraint cuts away a considerable area from the otherwise allowed region. Including a 10% error in theoretical calculations of the decay, the constraints from $b \rightarrow s\gamma$ loosen considerably, and all of the otherwise allowed WMAP area becomes available. The lightest Higgs is always lighter than 114 GeV, but heavier than 91 GeV.

The effect of varying the universal trilinear coupling A_0 is shown in Fig. 2. In contrast to the previous figure, here the sign of the μ -parameter is chosen to be negative. In general, giving the A_0 -parameter a nonzero value tends to increase the relic density. Increasing $|A_0|$ will help the $m_{H_u}^2$ to run larger negative values during the RG-




FIG. 1: Relic density $\Omega_\chi h^2$ in the representation **54** with $H = \text{SU}(2) \times \text{SO}(7)$ in the (M_2, m_0) plane for $\tan \beta = 10$, $\text{sgn}(\mu) = +1$, $A_0 = 0$. In **1a** the dark shaded areas represent the larger relic density. The filling denoted by wmap is the WMAP preferred region, lep shows an area, where the experimental mass limits are not met, rge shows an area where there is no radiative EWSB, and lsp the area where neutralino is not the LSP. h encloses the area with $m_h < 114$ GeV, and in the following figures bsg the area disallowed by $b \rightarrow s\gamma$ limits. In **1b** the NLSP within the same region is plotted, and in **1c** the $\tilde{\chi}_1^0$ composition.

evolution, and therefore to increase the actual (absolute) value for the μ -parameter via the rEWSB. This in turn favors the bino component in the $\tilde{\chi}_1^0$ composition in the $\text{SU}(2) \times \text{SO}(7)$ chain. The effect can be seen in the LSP-composition Fig. **2c**. The preferred WMAP region still follows the transition zone of the $\tilde{\chi}_1^0$ from bino to Higgsino. The effect of negative μ is most visible in the fact that the NLSP in this case is $\tilde{\chi}_2^0$, since negative μ tends to increase the lightest chargino mass (at least in the limit of $|M_2| < |\mu|$). The $b \rightarrow s\gamma$ constrains an area with negative A_0 values at $M_2 < 1$ TeV, and again including also the theoretical error, the constrained area becomes considerably smaller. The light Higgs boson mass is heavier than 114 GeV above the dash-dotted line.

In Fig. **3** the relic density, NLSP and $\tilde{\chi}_1^0$ composition are plotted for the $\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$ breaking chain. In this breaking chain the $|M_1|$ and $|M_2|$ values are closer to each other at the EW-scale than in the $\text{SU}(2) \times \text{SO}(7)$ chain, but still wider spread than in the singlet. That has an effect of increasing the μ value, therefore resulting in a smaller Higgsino component for the equal M_2 values for the two chains of the **54** dimensional representation. Again the WMAP allowed narrow region follows the transition from the $\tilde{\chi}_1^0$ from bino to Higgsino. The $b \rightarrow s\gamma$ and Higgs boson 114 GeV limits cut pieces from near the low M_2, m_0 -values. The NLSP is mostly the lightest chargino, but an interesting region exists with small m_0 near the area, where $\tilde{\chi}_1^0$ is no longer the LSP; along the line of transition from the stau NLSP to the smuon NLSP there is a narrow region, where $\tilde{\chi}_1^0$, stau and smuon masses are very close to each other, and the coannihilations may reduce the relic density to an acceptable level. For example, for $M_2 = 1400$ GeV, the LSP mass is

around 410 GeV, and for $M_2 = 1000$ GeV, the LSP mass is 290 GeV for the WMAP-preferred area. However, this area is highly prone to numerical subtleties, and the ordering of the LSP identity changes in the preferred relic density range, when comparing the output of different spectrum calculators. Reducing the m_0 parameter further makes the stau become the LSP. From the collider point of view, such regions may be especially interesting, as they would lead to quasistable smuons or staus.

2. Representation 210

In the representation **210** we inspected the breaking chain through the intermediate gauge group $\text{SU}(5) \times \text{U}(1)$, called flipped $\text{SU}(5)$ [29, 30, 31, 32]. In Fig. **4a** the area of preferred thermal relic density in the representation **210** is plotted for a set of (GUT scale) parameters. For the chosen parameters, rather large WMAP-preferred regions are found for large values of M_2 and/or m_0 parameters. When not Higgsino, the lightest neutralino is expected to be wino, rather than bino (see Table **II**), and therefore the neutralino relic density to be very small. In general, due to the wino being the smallest of the two electroweak gaugino parameters, it characterizes the lightest neutralino. Since the lightest chargino is characterized also by this parameter, for a large part of the parameter space, the masses of the $\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm$ are very close to each other, which boosts the rapid neutralino relic density annihilation. The situation resembles the one arising in the anomaly mediated SUSY breaking scenario, where also both the lightest neutralino and chargino are characterized by the wino mass param-




FIG. 2: Representation **54**: $H = \text{SU}(2) \times \text{SO}(7)$, $\tan \beta = 10$, $\text{sgn}(\mu) = -1$, $m_0 = 1$ TeV. Otherwise as in Fig. 1.




FIG. 3: Representation **54** with $H = \text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$ in the (M_2, m_0) plane for $\tan \beta = 10$, $\text{sgn}(\mu) = +1$, $A_0 = 0$. Otherwise as in Fig. 1.

eter.

Since both the wino and Higgsino have a large annihilation cross section, the WMAP-preferred relic density region does not have to follow the transition zone of $\tilde{\chi}_1^0$ from wino to Higgsino. The increase of the relic density to the observed level is mainly due to the increase of the mass parameters M_2 and m_0 . An interesting change in the pattern can be seen on the diagonal of the figures, where the $\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm$, and also at some point $\tilde{\chi}_2^0$, masses are very close to each other leading to enhanced coannihilation through the processes $\tilde{\chi}_1^0 \tilde{\chi}_1^\pm \rightarrow q_u \bar{q}_d$ and $\tilde{\chi}_1^0 \tilde{\chi}_1^0, \tilde{\chi}_1^\pm \tilde{\chi}_1^\pm \rightarrow q \bar{q}, \ell \bar{\ell}, W^+ W^-$, which allows the acceptable parameter space to extend to very large values of M_2 and m_0 .

The WMAP-preferred region is very wide as compared to, *e.g.*, the universal model or other models in $\text{SU}(5)$ [22, 33]. The spectrum is relatively heavy for the WMAP-preferred region, around a couple of TeV. This leads to

a wide range in the parameter space, but from the point of view of the coming Large Hadron Collider such a mass spectrum may be problematic. However, the representation **210** produces naturally a neutralino with mass around a TeV, which seems to be favorable in the view of the recent PAMELA [34] and ATIC [35] results of the excess positron and positron+electron flux. Under certain circumstances, a nearby clump of 600-1000 GeV neutralino LSP could fit into these observations [36].

In Fig. 5 the area of preferred thermal relic density in the representation **210** is plotted for the same set of (GUT scale) parameters as in Fig. 4, except that now the trilinear A_0 -parameter is varied along the y-axis, and the m_0 is set to 1 TeV. The interesting feature in this figure is the existence of the pseudoscalar Higgs annihilation channel through the M_2 -values. This reduces greatly the relic density in the parameter space where the LSP mass equals or is less than half of the A -Higgs mass.




FIG. 4: Representation **210**: $\tan\beta = 10$, $\text{sgn}(\mu) = +1$, $A_0 = 0$. Otherwise as in Fig. 1.




FIG. 5: Relic density in **Rep 210**: $\tan\beta = 10$, $\text{sgn}(\mu) = +1$, $m_0 = 1$ TeV. The collider constraints are fulfilled.

This has the effect of pushing the WMAP-preferred relic density region to heavier neutralino masses, and therefore to larger M_2 values. The width of the WMAP region is naturally the same as before, since the top of the figure, where $A_0 = 0$, coincides with Fig. 4 with $m_0 = 1000$ GeV. With these parameters, the NLSP is always the lighter chargino, and the lightest neutralino is dominantly a wino, which can also be read from Fig. 4c.

III. DISCUSSION AND SUMMARY

We studied the dark matter allowed regions in the SO(10) GUT representations, of which all but the singlet may lead to nonuniversal gaugino masses. The WMAP-preferred relic density regions are quite distinct for different representations, thus leading to quite different particle spectra for each representation.

In the representation **54**, the lightest neutralino is predominantly a bino, leading to the narrow areas of the WMAP-favored region. The excessive relic density is diluted either by increasing the Higgsino component or by coannihilation with other particles. The breaking chain $SU(2) \times SO(7)$ predicts an upper limit for the lighter chargino mass for the chosen parameters. For part of the WMAP allowed region, the whole neutralino and chargino spectrum is lighter than the spectrum of sfermions. For the $SU(4) \times SU(2) \times SU(2)$ breaking chain, the relic density area is narrow in the parameter space. Interestingly, there may exist a region, where the stau, smuon and the lightest neutralino masses are in a very close range to each other. This can lead to long-lived staus and smuons, which may be stable in the collider time scale.

In the **210** dimensional representation the lightest neutralino is either wino or Higgsino, which leads to a low thermal relic density. In addition, the lightest chargino and the lightest neutralino tend to be close in mass, thus providing a coannihilation channel. The preferred relic density area is quite large. The sparticle spectrum is heavy, as compared to the universal mSUGRA case. Only in a small part of the WMAP-preferred parameter region are a few SUSY particles expected to be within the kinematic reach of the LHC.

In this work we have studied each representation separately. It is obvious that if several representations affect simultaneously the composition of neutralinos, the possible WMAP-preferred region in the parameter space may be relaxed. However, if the neutralinos and charginos are found with a certain mass pattern, it helps to understand the relation of the lightest neutralino with dark matter, if characteristics of each representation are known.

IV. ACKNOWLEDGMENTS

The work of K.H. is supported by the Academy of Finland (Project No. 115032). The work of J.L. is supported

by the Bundesministerium für Bildung und Forschung, Berlin-Bonn.

-
- [1] D. N. Spergel et al. (WMAP), *Astrophys. J. Suppl.* **170**, 377 (2007), [astro-ph/0603449](#).
 - [2] K. Huitu, J. Laamanen, P. N. Pandita, and S. Roy, *Phys. Rev.* **D72**, 055013 (2005), [hep-ph/0502100](#).
 - [3] H. Fritzsch and P. Minkowski, *Ann. Phys.* **93**, 193 (1975).
 - [4] Z.-Y. Zhao, *J. Phys.* **G8**, 1019 (1982).
 - [5] M. Drees and J. M. Kim (2008), [0810.1875](#).
 - [6] S. Bhattacharya, A. Datta, and B. Mukhopadhyaya, *JHEP* **10**, 080 (2007), [0708.2427](#).
 - [7] B. Ananthanarayan and P. N. Pandita, *Int. J. Mod. Phys.* **A22**, 3229 (2007), [0706.2560](#).
 - [8] R. N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980).
 - [9] B. Bajc, G. Senjanovic, and F. Vissani, in **Budapest 2001, High energy physics* hep2001/198* (2001), [hep-ph/0110310](#).
 - [10] S. Dimopoulos and F. Wilczek, in *Proceedings of the 19th Course of the International School of Subnuclear Physics, Erice, Italy, 1981*, edited by A. Zichichi (Plenum Press, New York, 1983).
 - [11] E. Cremmer, S. Ferrara, L. Girardello, and A. Van Proeyen, *Phys. Lett.* **B116**, 231 (1982).
 - [12] S. Ferrara, L. Girardello, and H. P. Nilles, *Phys. Lett.* **B125**, 457 (1983).
 - [13] R. Slansky, *Phys. Rept.* **79**, 1 (1981).
 - [14] N. Chamoun, C.-S. Huang, C. Liu, and X.-H. Wu, *Nucl. Phys.* **B624**, 81 (2002), [hep-ph/0110332](#).
 - [15] G. F. Giudice and R. Rattazzi, *Nucl. Phys.* **B511**, 25 (1998), [hep-ph/9706540](#).
 - [16] C. S. Aulakh and S. K. Garg (2008), [0807.0917](#).
 - [17] C. S. Aulakh and S. K. Garg, *Nucl. Phys.* **B757**, 47 (2006), [hep-ph/0512224](#).
 - [18] B. Bajc, I. Dorsner, and M. Nemevsek, *JHEP* **11**, 007 (2008), [0809.1069](#).
 - [19] C. S. Aulakh and R. N. Mohapatra, *Phys. Rev.* **D28**, 217 (1983).
 - [20] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac, and N. Okada, *J. Math. Phys.* **46**, 033505 (2005), [hep-ph/0405300](#).
 - [21] S. F. King, J. P. Roberts, and D. P. Roy (2007), [arXiv:0705.4219](#).
 - [22] K. Huitu et al., *Eur. Phys. J.* **C58**, 591 (2008), [0808.3094](#).
 - [23] B. C. Allanach, *Comput. Phys. Commun.* **143**, 305 (2002), [hep-ph/0104145](#).
 - [24] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput. Phys. Commun.* **149**, 103 (2002), [hep-ph/0112278](#).
 - [25] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput. Phys. Commun.* **174**, 577 (2006), [hep-ph/0405253](#).
 - [26] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput. Phys. Commun.* **176**, 367 (2007), [hep-ph/0607059](#).
 - [27] E. Barberio et al. (Heavy Flavor Averaging Group (HFAG)) (2007), [0704.3575](#).
 - [28] *Search for charged Higgs bosons: Preliminary combined results using LEP data collected at energies up to 209-GeV* (2001), [hep-ex/0107031](#), [hep-ex/0107031](#).
 - [29] A. De Rujula, H. Georgi, and S. L. Glashow, *Phys. Rev. Lett.* **45**, 413 (1980).
 - [30] S. M. Barr, *Phys. Lett.* **B112**, 219 (1982).
 - [31] J. P. Derendinger, J. E. Kim, and D. V. Nanopoulos, *Phys. Lett.* **B139**, 170 (1984).
 - [32] I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, *Phys. Lett.* **B194**, 231 (1987).
 - [33] K. Huitu, J. Laamanen, and S. Roy (2007), [0710.2058](#).
 - [34] O. Adriani et al. (2008), [0810.4994](#).
 - [35] J. Chang et al., *Nature* **456**, 362 (2008).
 - [36] D. Hooper, A. Stebbins, and K. M. Zurek (2008), [0812.3202](#).